



## **About MACUGEN® (pegaptanib sodium injection)**

In December 2004, MACUGEN became the first anti-VEGF inhibitor to be FDA approved for the treatment of wet AMD. MACUGEN is effective at treating all types of wet AMD by slowing vision loss. It is available by prescription only.

While MACUGEN is effective at slowing vision loss, it is still important to detect wet AMD as early as possible to help preserve vision. Wet AMD is a serious and progressive condition that can cause rapid central vision loss. Without treatment, patients can have severe vision loss in the affected eye within months of diagnosis.

Learn more about the SYMPTOMS OF WET AMD

## **How MACUGEN Works**

MACUGEN works differently from all other treatments for wet AMD. MACUGEN selectively blocks an essential signal in the eye that is believed to cause abnormal blood vessels to grow and leak. By helping to slow the growth of these abnormal blood vessels, MACUGEN can slow the damage they do to your sight. MACUGEN is not a cure, but it can help to slow further vision loss and help preserve what vision you currently have.

Ask your eye doctor how MACUGEN can help you. If you don't have an eye doctor, use the Eye Doctor Finder to locate one near you.

Learn more about MACUGEN and wet AMD by downloading our educational brochure.

You will need Adobe® Reader® to download this file. If you don't have Adobe Reader, you can download it for free.

MACUGEN is administered into the eye every 6 weeks. Your doctor will determine how long you'll need treatment. Be sure to attend follow-up appointments for eye exams and MACUGEN treatments according to your doctor's recommendations.

## **How MACUGEN® (pegaptanib sodium injection) is administered**

MACUGEN is not a pill; no medicines in a pill form have been approved to treat wet AMD. MACUGEN is delivered by injection directly into the eye to reach the back of the eye where it is needed. Your eye doctor is trained in treating eye diseases and in giving eye injections. The needle used is very small. The needle is inserted near the edge of the eye, not in the center. While administering MACUGEN, your eye doctor will ask you to look in the opposite direction, so you will not see the needle during the injection procedure. During your treatment there are additional procedures your eye doctor will complete to administer MACUGEN, such as numbing the surface of your eye.



MACUGEN is generally well tolerated. In clinical studies, the vast majority of patients continued their injections of MACUGEN every 6 weeks for 2 years, and over 75,000 people have now received MACUGEN. If you have specific questions about the injection procedure, ask your treating eye doctor.

## Learn more about TREATMENT RESULTS WITH MACUGEN

To listen to the full prescribing information regarding MACUGEN, please click on the link entitled "Important Safety Information" on the menu bar located on the left of the screen.

MACUGEN is approved to treat wet age-related macular degeneration. It is available by prescription only.

Not all people respond the same to MACUGEN, so individual results may vary. The safety or efficacy of MACUGEN has not been proven beyond 2 years.

### **Important safety information**

With MACUGEN, you may have some side effects, mostly in the eye and due to the injection procedure. The most common side effects are burning sensation, eye pain, redness, light sensitivity, vision loss, blurred vision, visual disturbances, high blood pressure, and cataract. For a complete list of side effects, please ask your doctor or see the prescribing information.

You should not receive MACUGEN if you have an infection in or around your eye or if you are hypersensitive (allergic) to pegaptanib sodium or any of the other ingredients.

A serious eye infection can sometimes develop after an injection into the eye. Signs of a serious infection may include eye pain, light sensitivity, and/or vision changes. Check with a retinal physician immediately if you experience any of these symptoms, so you can be treated early if an infection occurs.

Eye injections like those with MACUGEN can increase eye pressure. This is something you would not notice; however, your retinal physician may do some extra tests after your injection to make sure there are no complications.

In rare cases, allergic reactions have been known to occur. Tell your retinal physician about any known allergies.

Please follow your retinal physician's recommendations for your therapy. If you have any questions about MACUGEN, ask your retinal physician.

The health information contained herein is provided for educational purposes only and is not intended to replace discussions with a health care provider. All decisions



IT TAKES ALL TYPES™



regarding patient care must be made with a health care provider, considering the unique characteristics of the patient.

Copyright 2006 (OSI) Eyetech Pharmaceuticals, Inc., and Pfizer Inc. All rights reserved.

The product information in this site is intended only for residents of the United States. The products discussed herein may have different product labeling in different countries.

Pfizer is a licensee of the TRUSTe Privacy Program.  
MU277755