



# MACUGEN® (pegaptanib sodium injection)




## DESCRIPTION


MACUGEN® (pegaptanib sodium injection) is a sterile, aqueous solution containing pegaptanib sodium for intravitreal injection. Macugen is supplied in a single-dose, pre-filled syringe and is formulated as a 3.47 mg/mL solution, measured as the free acid form of the oligonucleotide. The active ingredient is 0.3 mg of the free acid form of the oligonucleotide without polyethylene glycol, in a nominal volume of 90 µL. This dose is equivalent to 1.6 mg of pegaptanib sodium (pegylated oligonucleotide) or 0.32 mg when expressed as the sodium salt form of the oligonucleotide moiety. The product is a sterile, clear, preservative-free solution containing sodium chloride, monobasic sodium phosphate monohydrate, dibasic sodium phosphate heptahydrate, hydrochloric acid, and/or sodium hydroxide to adjust the pH and water for injection.

Pegaptanib sodium is a covalent conjugate of an oligonucleotide of twenty-eight nucleotides in length that terminates in a penylamino linker, to which two 20-kilodalton monomethoxy polyethylene glycol (PEG) units are covalently attached via the two amino groups on a lysine residue.

Pegaptanib sodium is represented by the following structural formula:



Where R is



and n is approximately 450.

The chemical name for pegaptanib sodium is as follows: RNA, ((2'-deoxy-2'-fluoro)C-G<sub>m</sub>-G<sub>m</sub>-A-A-(2'-deoxy-2'-fluoro)U-(2'-deoxy-2'-fluoro)C-A<sub>m</sub>-G<sub>m</sub>-(2'-deoxy-2'-fluoro)U-(2'-deoxy-2'-fluoro)U-A<sub>m</sub>-(2'-deoxy-2'-fluoro)U-A<sub>m</sub>-(2'-deoxy-2'-fluoro)U-G<sub>m</sub>-(2'-deoxy-2'-fluoro)U-G<sub>m</sub>-(2'-deoxy-2'-fluoro)U-(2'-deoxy-2'-fluoro)U-A<sub>m</sub>-(2'-deoxy-2'-fluoro)U-A<sub>m</sub>-(2'-deoxy-2'-fluoro)C-A<sub>m</sub>-(2'-deoxy-2'-fluoro)U-(2'-deoxy-2'-fluoro)C-(2'-deoxy-2'-fluoro)C-G<sub>m</sub>-(3'→3')dT, 5'-ester with α,α'-[4,12-dioxo-6-[[[5-(phosphonoxy)pentyl]amino]carbonyl]-3,13-dioxo-5,11-diaza-1,15-pentadecanediyl]bis[ω-methoxypropyl(oxy-1,2-ethanediyl)], sodium salt.

The molecular formula for pegaptanib sodium is C<sub>294</sub>H<sub>342</sub>F<sub>13</sub>N<sub>107</sub>Na<sub>28</sub>O<sub>188</sub>P<sub>28</sub>[C<sub>2</sub>H<sub>4</sub>O]<sub>n</sub> (where n is approximately 900) and the molecular weight is approximately 50 kilodaltons. Macugen is formulated to have an osmolality of 280-360 mOsm/Kg, and a pH of 6-7.

## CLINICAL PHARMACOLOGY

### Mechanism of Action

Pegaptanib is a selective vascular endothelial growth factor (VEGF) antagonist. VEGF is a secreted protein that selectively binds and activates its receptors located primarily on the surface of vascular endothelial cells. VEGF induces angiogenesis, and increases vascular permeability and inflammation, all of which are thought to contribute to the progression of the neovascular (wet) form of age-related macular degeneration (AMD), a leading cause of blindness. VEGF has been implicated in blood retinal barrier breakdown and pathological ocular neovascularization.

Pegaptanib is an aptamer; a pegylated modified oligonucleotide, which adopts a three-dimensional conformation that enables it to bind to extracellular VEGF. Under in vitro testing conditions, pegaptanib binds to the major pathological VEGF isoform, extracellular VEGF<sub>165</sub>, thereby inhibiting VEGF<sub>165</sub> binding to its VEGF receptors. The inhibition of VEGF<sub>165</sub>, the rodent counterpart of human VEGF<sub>165</sub>, was effective at suppressing pathological neovascularization.

### Pharmacokinetics

#### Absorption

In animals, pegaptanib is slowly absorbed into the systemic circulation from the eye after intravitreal administration. The rate of absorption from the eye is the rate limiting step in the disposition of pegaptanib in animals and is likely to be the rate limiting step in humans.

In humans, a mean maximum plasma concentration of about 80 ng/mL occurs within 1 to 4 days after a 3 mg monocular dose (10 times the recommended dose). The mean area under the plasma concentration-time curve (AUC) is about 25 µg·hr/mL at this dose.

## Distribution/Metabolism/Excretion

Twenty-four hours after intravitreal administration of a radiolabeled dose of pegaptanib to both eyes of rabbits, radioactivity was mainly distributed in vitreous fluid, retina, and aqueous fluid. After intravitreal and intravenous administrations of radiolabeled pegaptanib to rabbits, the highest concentrations of radioactivity (excluding the eye for the intravitreal dose) were obtained in the kidney. In rabbits, the component nucleotide, 2'-fluorouridine is found in plasma and urine after single radiolabeled pegaptanib intravenous and intravitreal doses. In rabbits, pegaptanib is eliminated as parent drug and metabolites primarily in the urine. Based on preclinical data, pegaptanib is metabolized by endo- and exonucleases.

In humans, after a 3 mg monocular dose (10 times the recommended dose), the average (± standard deviation) apparent plasma half-life of pegaptanib is 10 (±4) days.

## Special Populations

Plasma concentrations do not appear to be affected by age or gender, but have not been studied in patients under the age of 50.

## Renal Insufficiency

Dose adjustment for patients with renal impairment is not needed when administering the 0.3 mg dose.

Following a single 3 mg dose (10 times the recommended dose), in patients with severe (N=7), moderate (N=18), and mild (N=10) renal impairment, the mean (CV%) pegaptanib AUC values were 37.8 (17%), 26.7 (31%), and 23.6 (21%) µg·hr/mL, respectively. The corresponding C<sub>max</sub> values were 96.8 (23%), 81.6 (29.2%), and 66.5 (47%) ng/mL, respectively.

In patients with renal impairment, following administration of 3 mg pegaptanib doses every 6 weeks, the last detectable pegaptanib concentrations in plasma after the fourth dose were highly variable (ranging from 8 ng/mL to 66 ng/mL) and the variability was more pronounced in patients with severe renal impairment.

## Hemodialysis

Macugen has not been studied in patients requiring hemodialysis.

## Hepatic Impairment

Macugen has not been studied in patients with hepatic impairment.


## Clinical Studies

Macugen was studied in two controlled, double-masked, and identically designed randomized studies in patients with neovascular AMD. Patients were randomized to receive control (sham treatment) or 0.3 mg, 1 mg or 3 mg Macugen administered as intravitreal injections every 6 weeks for 48 weeks. A total of approximately 1200 patients were enrolled with 892 patients receiving Macugen and 298 receiving a sham injection. The median age of the patients was 77 years. Patients received a mean 8.5 treatments out of a possible 9 total treatments across all treatment arms. Patients were re-randomized between treatment and no treatment during the second year. Patients who continued treatment in year 2 received a mean of 16 treatments out of a possible total 17 overall.

The two trials enrolled patients with neovascular AMD characteristics including classic, occult, and mixed lesions of up to 12 disc areas and baseline visual acuity in the study eye between 20/40 and 20/320. The primary efficacy endpoint was the proportion of patients losing less than 15 letters of visual acuity, from baseline up to 54 week assessment. Verteporfin photodynamic therapy (PDT) usage was permitted at the discretion of the investigators in patients with predominantly classic lesions.


The groups treated with Macugen 0.3 mg exhibited a statistically significant result in both trials for the primary efficacy endpoint at 1 year: Study EOP1003, Macugen 73% vs. Sham 60%; Study EOP1004, Macugen 67% vs. Sham 53%. Concomitant use of PDT overall was low. More sham treated patients (75/296) received PDT than Macugen 0.3 mg treated patients (58/294). On average, Macugen 0.3 mg treated patients and sham treated patients continued to experience vision loss. However, the rate of vision decline in the Macugen treated group was slower than the rate in the patients who received sham treatment. See Figure 1.

Figure 1



At the end of the first year (week 54), approximately 1050 of the original 1200 patients were re-randomized to either continue the same treatment or to discontinue treatment through week 102. See Figure 2. Macugen was less effective during the second year than during the first year. The percentage of patients losing less than 15 letters from baseline to week 102 was: Study EOP1003, Macugen 38/67 (57%); Sham 30/54 (56%); Study EOP1004, Macugen 40/66 (61%); Sham 18/53 (34%).

Figure 2



Dose levels above 0.3 mg did not demonstrate any additional benefit.

The safety or efficacy of Macugen beyond 2 years has not been demonstrated.

## INDICATIONS AND USAGE

Macugen is indicated for the treatment of neovascular (wet) age-related macular degeneration.

## CONTRAINDICATIONS

Macugen is contraindicated in patients with ocular or periorcular infections.

Macugen is contraindicated in patients with known hypersensitivity to pegaptanib sodium or any other excipient in this product.

## WARNINGS

Intravitreal injections including those with Macugen have been associated with endophthalmitis. Proper aseptic injection technique should always be utilized when administering Macugen. In addition, patients should be monitored during the week following the injection to permit early treatment, should an infection occur (see **DOSAGE AND ADMINISTRATION**).

Increases in intraocular pressure have been seen within 30 minutes of injection with Macugen. Therefore, intraocular pressure as well as the perfusion of the optic nerve head should be monitored and managed appropriately.

## PRECAUTIONS

### General

FOR OPHTHALMIC INTRAVITREAL INJECTION ONLY.

Rare cases of anaphylaxis/anaphylactoid reactions, including angioedema, have been reported in the post-marketing experience following the Macugen intravitreal administration procedure (see **ADVERSE EVENTS** and **DOSAGE AND ADMINISTRATION**).

### Information for Patients

In the days following Macugen administration, patients are at risk for the development of endophthalmitis. If the eye becomes red, sensitive to light, painful or develops a change in vision, the patient should seek the immediate care with their ophthalmologist.

### Drug Interactions

Drug interaction studies have not been conducted with Macugen. Pegaptanib is metabolized by nucleases and is generally not affected by the cytochrome P450 system.



Two early clinical studies conducted in patients who received Macugen alone and in combination with PDT revealed no apparent difference in the plasma pharmacokinetics of pegaptanib.

#### Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenicity studies with pegaptanib have not been conducted.

Pegaptanib and its monomer component nucleotides (2'-MA, 2'-MG, 2'-FU, 2'-FC) were evaluated for genotoxicity in a battery of *in vitro* and *in vivo* assay systems. Pegaptanib, 2'-O-methyladenosine (2'-MA), and 2'-O-methylguanosine (2'-MG) were negative in all assay systems evaluated. 2'-fluorouridine (2'-FU) and 2'-fluorocytidine (2'-FC) were nonclastogenic and were negative in all *S. typhimurium* tester strains, but produced a non-dose related increase in revertant frequency in a single *E. coli* tester strain. Pegaptanib, 2'-FU, and 2'-FC tested negative in cell transformation assays.

No data are available to evaluate male or female mating or fertility indices.

#### Pregnancy

Teratogenic Effects: Pregnancy Category B.

Pegaptanib produced no maternal toxicity and no evidence of teratogenicity or fetal mortality in mice at intravenous doses of up to 40 mg/kg/day (about 7,000 times the recommended human monocular ophthalmic dose of 0.3 mg/eye). Pegaptanib crosses the placenta in mice.

There are no studies in pregnant women. The potential risk to humans is unknown. Macugen should be used during pregnancy only if the potential benefit to the mother justifies the potential risk to the fetus.

#### Nursing Mothers

It is not known whether pegaptanib is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when Macugen is administered to a nursing woman.

#### Pediatric Use

Safety and effectiveness of Macugen in pediatric patients have not been studied.

#### Geriatric Use

Approximately 94% (834/892) of the patients treated with Macugen were  $\geq$  65 years of age and approximately 62% (553/892) were  $\geq$  75 years of age. No difference in treatment effect or systemic exposure was seen with increasing age.

#### **ADVERSE EVENTS**

Serious adverse events related to the injection procedure occurring in  $<$  1% of intravitreal injections included endophthalmitis (see **WARNINGS**), retinal detachment, and iatrogenic traumatic cataract.

The most frequently reported adverse events in patients treated with Macugen 0.3 mg for up to two years were anterior chamber inflammation, blurred vision, cataract, conjunctival hemorrhage, corneal edema, eye discharge, eye irritation, eye pain, hypertension, increased intraocular pressure (IOP), ocular discomfort, punctate keratitis, reduced visual acuity, visual disturbance, vitreous floaters, and vitreous opacities. These events occurred in approximately 10-40% of patients.

The following events were reported in 6-10% of patients receiving Macugen 0.3 mg therapy:

Ocular: blepharitis, conjunctivitis, photopsia, vitreous disorder.

Non-Ocular: bronchitis, diarrhea, dizziness, headache, nausea, urinary tract infection.

The following events were reported in 1-5% of patients receiving Macugen 0.3 mg therapy:

Ocular: allergic conjunctivitis, conjunctival edema, corneal abrasion, corneal deposits, corneal epithelium disorder, endophthalmitis, eye inflammation, eye swelling, eyelid irritation, meibomianitis, mydriasis, periorbital hematoma, retinal edema, vitreous hemorrhage.

Non-Ocular: arthritis, bone spur, carotid artery occlusion, cerebrovascular accident, chest pain, contact dermatitis, contusion, diabetes mellitus, dyspepsia, hearing loss, pleural effusion, transient ischemic attack, urinary retention, vertigo, vomiting.

Post-Marketing Experience: Anaphylaxis/anaphylactoid reactions, including angioedema, have been identified during postapproval use of Macugen. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure (see **PRECAUTIONS** and **DOSAGE AND ADMINISTRATION**).

#### **OVERDOSAGE**

Doses of Macugen up to 10 times the recommended dosage of 0.3 mg have been studied. No additional adverse events have been noted but there is decreased efficacy with doses above 1 mg.

#### **DOSAGE AND ADMINISTRATION**

Macugen 0.3 mg should be administered once every six weeks by intravitreal injection into the eye to be treated.

Macugen should be inspected visually for particulate matter and discoloration prior to administration.


Administration of the syringe contents involves assembly of the syringe with the administration needle. The injection procedure should be carried out under controlled aseptic conditions, which includes the use of sterile gloves, a sterile drape, and a sterile eyelid speculum (or equivalent). When ready to assemble syringe and administer injection, carefully peel open pouches, remove contents, and place on sterile field. If upon opening the pouch, the plastic clip is missing or not attached to the syringe, the syringe should not be used.

**To avoid compromising the sterility of the product, do not pull back on the plunger.**

1. Remove the syringe from the plastic clip.
2. Twist off cap.
3. Attach the sterile BD<sup>®</sup> 30G 1/2" Precision Glide<sup>®</sup> administration needle (included) to the syringe by screwing it into the syringe tip. --Another sterile administration needle may be used in lieu of the one included. Remove the plastic needle shield from the needle.
4. Holding the syringe with the needle pointing up, check the syringe for bubbles. If there are bubbles, gently tap the syringe with your finger until the bubbles rise to the top of the syringe. **SLOWLY** depress the plunger to eliminate all the bubbles and to expel the excess drug so that the **top edge of the 3rd rib on the plunger stopper aligns with the pre-printed black dosing line** (See Fig 2, below).
5. Inject the entire contents of the syringe.

#### **PRIOR to Injection**


**Fig 1. Before** expelling air bubble and excess drug



(Actual air bubble formation may vary)

#### **READY for Injection**

**Fig 2. After** expelling air bubble and excess drug



The patient's medical history for hypersensitivity reactions should be evaluated prior to performing the intravitreal procedure (see **PRECAUTIONS** and **ADVERSE EVENTS**). Adequate anesthesia and a broad-spectrum microbicide should be given prior to the injection.

Following the injection, patients should be monitored for elevation in intraocular pressure and for endophthalmitis. Monitoring may consist of a check for perfusion of the optic nerve head immediately after the injection, tonometry within 30 minutes following the injection, and biomicroscopy between two and seven days following the injection. Patients should be instructed to report any symptoms suggestive of endophthalmitis without delay.

No special dosage modification is required for any of the populations that have been studied (i.e. gender, elderly).

The safety and efficacy of Macugen therapy administered to both eyes concurrently have not been studied.

#### **HOW SUPPLIED**

Macugen (pegaptanib sodium injection) is supplied in a sterile foil pouch as a single-use glass syringe pre-filled with 0.3 mg of Macugen<sup>®</sup> in a nominal 90  $\mu$ L deliverable volume pack. A sterile packaged BD<sup>®</sup> single use 30G x 1/2" Precision Glide<sup>®</sup> Luer Lok<sup>®</sup> needle is supplied in a separate pouch. The foil pouch and needle are packaged together in a carton.

#### Storage

Store in the refrigerator at 2° to 8°C (36° to 46°F). Do not freeze or shake vigorously.

Rx only.

BD and Precision Glide Luer Lok<sup>®</sup> are registered trademarks of Becton Dickinson & CO, Franklin Lakes, New Jersey 07417

NDC 68782-001-02

Manufactured by:  
Gilead Sciences, Inc  
650 Cliffs Drive  
San Dimas, CA 91773

For:  
OSI Eyetech, Inc.  
140 East Hanover Avenue  
Cedar Knolls, NJ 07927

70-6127-00-2

Revised 4/2008

